

# Thyroid Hormones Interact with Glucocorticoids to Affect Somatotroph Abundance in Chicken Embryonic Pituitary Cells *in Vitro*

LIXIN LIU, CARLTON E. DEAN, AND TOM E. PORTER

*Department of Animal and Avian Sciences (L.L., C.E.D., T.E.P.) and Molecular and Cell Biology Program (T.E.P.), University of Maryland, College Park, Maryland 20742*

Our laboratory has reported that somatotroph differentiation occurs between d 14 and d 16 of chicken embryonic development and that corticosterone (CORT) can induce somatotroph differentiation at an earlier age *in vitro* and *in vivo*. The objective of the present study was to test for thyroid hormone-CORT interactions on somatotroph differentiation *in vitro*. Pituitary cells from d 11 chicken embryos were treated with CORT and thyroid hormones, and GH-producing somatotrophs were detected by reverse hemolytic plaque assays and immunocytochemistry. We found that thyroid hormones can act synergistically with CORT to further augment the abundance of somatotrophs *in vitro* but have little to no effect on their own. Both T<sub>4</sub> and T<sub>3</sub> could act synergistically with CORT to increase somatotroph abundance, but the effects of

T<sub>3</sub> were biphasic, inhibiting CORT actions at higher concentrations. The monodeiodination inhibitor iopanoic acid inhibited the synergistic effect of T<sub>4</sub> on CORT induction of GH cells *in vitro* but not the synergistic effect of CORT and T<sub>3</sub> or the effect of CORT alone. Furthermore, T<sub>3</sub> treatment overcame the iopanoic acid-induced reduction in the T<sub>4</sub>-CORT effect. Our findings indicate that thyroid hormones act synergistically with CORT to further augment the abundance of somatotrophs *in vitro* and that conversion of T<sub>4</sub> to T<sub>3</sub> within the pituitary is involved in T<sub>4</sub> modulation of somatotroph abundance. Somatotroph differentiation during normal development may be regulated by complex interactions of hormones produced by the embryonic thyroid and adrenal glands. (*Endocrinology* 144: 3836–3841, 2003)

THE DEVELOPMENT OF the pituitary gland has been the subject of many studies aimed at elucidating the molecular mechanisms of generating diverse cell phenotypes from a common precursor within an organ (1, 2). The differentiation of pituitary cells is not autonomous but is controlled by intracellular and extracellular signals. The anterior pituitary-specific transcription factor Pit-1/GHF-1 is expressed in the thyrotroph, somatotroph, and lactotroph and is required for the proliferation and maintenance of these three cell types as well as the transcriptional activation of their corresponding trophic hormone genes (3). However, its expression pattern indicates that Pit-1 protein is not sufficient for somatotroph differentiation because GH is not expressed in the lactotroph or thyrotroph lineages, which also express Pit-1 protein. Thus, unknown cell-specific factors, in addition to Pit-1, exist that need to be identified to fully understand somatotroph differentiation during development.

Our laboratory is studying the mechanisms regulating differentiation of GH-secreting cells in the chicken anterior pituitary. During chicken embryonic development, which lasts 21 d, somatotrophs are rare before embryonic day (e) 12, and significant somatotroph differentiation occurs between e14 and e16 (4). GH-secreting cells do not differentiate spontaneously without an extrapituitary signal, and serum from e16 embryos but not e12 can induce somatotroph differentiation in culture (5). The adrenal glucocorticoid, corticosterone, is the active compound responsible for cell-differentiating activity of e16 chicken serum (6). Glucocorticoids

were found to induce somatotrophs in chicken embryos and fetal rats *in vitro* and *in vivo* (6–9). Furthermore, glucocorticoids were found to increase pituitary GH mRNA levels in rats (8), enhance GH mRNA stability in humans (10), and increase the level of GH gene expression and the number of cells that expressed GH mRNA in chickens (11).

Thyroid hormones play an important role in embryonic and neonatal development in mammals and birds. In humans, hypothyroidism severely impairs postnatal growth and reduces spontaneous nocturnal GH secretion (12, 13). T<sub>3</sub> was observed to increase GH mRNA accumulation in a cell line (GX) derived from a pituitary tumor of a boy with gigantism (14). In rats, thyroid hormones have been well documented to control GH synthesis. Thyroid hormones stimulated rat GH gene expression *in vivo* (15) and in several GH-producing pituitary cell lines (e.g. GH<sub>1</sub>, GH<sub>3</sub>, and GC) (16–18). In GC cells, T<sub>3</sub> was shown to rapidly stimulate GH gene transcription, and this effect was mediated by thyroid hormone receptors (19). However, thyroid hormones were found to exert their stimulatory action on fetal GH gene expression and GH cell abundance only in the presence of glucocorticoids (20). In chickens, the regulation of GH secretion by thyroid hormones is complex. The importance of T<sub>3</sub> to the normal growth process was demonstrated by the ability of exogenous T<sub>3</sub> to correct the growth deficit of hypophysectomized chickens (21). In contrast, administration of thyroid hormones to posthatch chickens decreased GH synthesis and secretion (22). However, nothing is known about the effects of thyroid hormones on somatotroph differentiation during chicken embryonic development.

Previous studies have evaluated the effects of combined

Abbreviations: e, Embryonic day; ICC, immunocytochemistry; IOP, iopanoic acid; RHPA, reverse hemolytic plaque assay.

thyroid hormone and glucocorticoid treatment on GH expression in the pituitary glands of fetal rats. Hemming *et al.* (23) found that  $T_3$ , which was individually ineffective, acted synergistically with cortisol to increase somatotroph density in pituitary tissue from d 14 rat fetuses cultured for 7 d. Because GH immunostaining in histology sections of the cultured tissue was evaluated in this study, qualitative rather than quantitative results were obtained. Nogami *et al.* (20) found that adding dexamethasone to the drinking water of pregnant rats induced GH and GH mRNA accumulation in the d 18 fetal pituitary gland. Additional ip  $T_4$  injections enhanced the effect of dexamethasone, whereas  $T_4$  exhibited no effect when given alone. Because treatments were administered to the dams rather than directly to the fetuses in these studies, the potential involvement of a treatment-induced maternal mediator of these effects could not be eliminated. This group also examined the effect of glucocorticoids and  $T_3$  on GH mRNA levels in cultured fetal pituitary glands (8). However, the abundance of somatotrophs was not evaluated in that study.

The purpose of the current study was to quantify direct effects of thyroid hormones and corticosterone on somatotroph abundance in cultures of chicken embryonic pituitary cells. Reverse hemolytic plaque assays (RHPA) and immunocytochemistry (ICC) were conducted to quantify the effects of these treatments on somatotroph abundance. The requirement for monodeiodination of  $T_4$  in induction of somatotroph differentiation was also tested.

## Materials and Methods

### Animals and pituitary dispersions

All animals used in the present study were Avian X Avian broiler strain chicken embryos purchased from Allen's Hatchery (Seaford, DE). All procedures with chicken embryos were approved by the Institutional Animal Care and Use Committee on this campus. Unless stated otherwise, all cell culture reagents were obtained from Life Technologies, Inc. (Grand Island, NY), and hormones and other chemicals were purchased from Sigma Chemical Co. (St. Louis, MO). All media were supplemented with 0.1% BSA, 100 U/ml penicillin G, and 100  $\mu$ g/ml streptomycin sulfate. Fertile eggs were placed in a humidified incubator (G.Q.F. Manufacturing, Savannah, GA) at 37.5 C. The typical length of incubation for chickens is 21 d. Embryos were removed on e11, and their anterior pituitary glands were isolated with the aid of a dissecting microscope. The pituitaries were then dissociated by a combination of trypsin digestion and mechanical agitation as described previously (4–7). The viability of the cells was assessed by the trypan blue dye exclusion method and was consistently greater than 95%.

### Extended cell cultures

Anterior pituitary cells were cultured according to the procedure described previously (4–6). Cells were plated ( $2.0 \times 10^5$  cells/well) in poly-L-lysine-coated 12-well tissue culture plates and allowed to attach for 45 min in 75  $\mu$ l DMEM. Wells were then filled with 2 ml serum-free medium (1:1 mixture of phenol red-free medium 199 and Ham's F-12, supplemented with 0.1% BSA, 5  $\mu$ g/ml human transferrin, 5  $\mu$ g/ml bovine insulin, 100 U/ml penicillin G, and 100  $\mu$ g/ml streptomycin sulfate) alone or medium containing different treatments. After culture for 3 or 6 d in a humidified incubator (37.5 C; 95% air-5%  $CO_2$ ), cells were harvested for detection of GH-secreting cells by RHPA or GH-containing cells by ICC.

### RHPAs

The assays were performed according to the protocol described in detail previously (4–6). Briefly, the harvested pituitary cells ( $1.0 \times 10^5$

/ml) were mixed with an equal volume of an 18% suspension of protein A-coated ovine erythrocytes and infused by capillary action into poly-L-lysine coated Cunningham chambers. After cells were allowed to attach for 45 min (37.5 C; 95% air-5%  $CO_2$ ), chambers were rinsed with DMEM to remove unattached cells. DMEM containing GH antiserum (1:40) and human GHRH ( $10^{-7}$  M) was then added to the resulting monolayer of cells, and replicate chambers were incubated for 20 h (three chambers per treatment). Plaque formation was subsequently induced by a 45-min incubation with guinea pig complement (1:40, in DMEM). The cells were then fixed with 2% glutaraldehyde in 0.9% saline and stained with methyl green. Chambers were analyzed using a light microscope. Data shown are the percentage of all pituitary cells that formed plaques (*i.e.* secreted GH), with at least 200 pituitary cells counted/assay chamber.

### ICC

The harvested pituitary cells were diluted in DMEM to a concentration of  $1.0 \times 10^4$  cells/ml and were then attached to poly-L-lysine coated 24-well tissue culture plates during a 1-h incubation period (37.5 C; 95% air-5%  $CO_2$ ). The plated cells were fixed with 3.7% formaldehyde in PBS for 20 min. Cells were permeabilized with 0.1% Tween 20/0.1% Triton X-100 (8 min), quenched with 0.3%  $H_2O_2$  (5 min), blocked with 1% normal goat serum (15 min), and incubated overnight with rabbit antichick GH (1:4000 in PBS) using the antiserum described above for the RHPA. The cells were rinsed with PBS and incubated with biotinylated goat antirabbit IgG for 30 min. The cells were further processed using rabbit avidin biotin complex kits according to the directions supplied by the manufacturer (Vector Laboratories, Burlingame, CA). VIP reagent (Vector) was used as substrate for the peroxidase. GH-containing cells were identified using an inverted light microscope, and data shown are the percentage of the total pituitary cell population that contained GH.

### Statistical analysis

The data reported are the mean  $\pm$  SEM from at least four completely separate experiments, with the number of replicate experiments provided in the legend to each figure. All data were analyzed by ANOVA using the mixed models procedure of SAS (SAS Institute, Cary, NC). Differences between treatments were tested using Tukey's Studentized range test and were considered significant at  $P < 0.05$ .

## Results

### Effect of corticosterone and $T_3$ alone or in combination on somatotroph abundance

We reported that corticosterone induces somatotroph differentiation within 2 d in cultures of chick embryo pituitary cells (6). We also showed that corticosterone and GHRH act synergistically after 3 d to further increase GH cell abundance (24). To extend our investigation of hormone interactions, pituitary cells from e11 chicks were cultured for 3 or 6 d with corticosterone ( $10^{-9}$  M) or  $T_3$  ( $10^{-11}$  to  $10^{-8}$  M) alone or in combination, and then cells were subjected to GH RHPA. Figure 1 presents the results from these trials. Corticosterone alone increased proportions of GH-secreting cells by 3 and 6 d to  $5.5 \pm 1.7$  (mean  $\pm$  pooled SE of the mean) and  $6.1 \pm 1.4\%$  vs. basal levels of  $1.2 \pm 0.6\%$  and  $0.3 \pm 0.2\%$ , respectively. No significant effects were found for any concentration of  $T_3$  alone. Therefore,  $T_3$  alone had no effect on the abundance of somatotrophs in cultures of chick embryonic pituitary cells. However, low doses of  $T_3$  in combination with corticosterone further increased the proportions of GH-secreting cells above that with corticosterone alone (to  $10.3 \pm 1.8\%$  at  $10^{-9}$  M  $T_3$  after 3 d and  $9.0 \pm 1.5\%$  at  $10^{-10}$  M  $T_3$  after 6 d). In contrast, high levels of  $T_3$  inhibited the stimulatory effect of corticosterone on proportions of GH-secreting cells




FIG. 1. The effect of corticosterone and T<sub>3</sub> on somatotroph abundance determined by RHPA. Pituitary cells from e11 chicks were cultured for 3 d (upper panel) or 6 d (lower panel) with corticosterone (10<sup>-9</sup> M) or T<sub>3</sub> (10<sup>-11</sup> to 10<sup>-8</sup> M) alone or in combination. Cells were harvested and subjected to RHPA for GH. Slides were incubated for 20 h in the presence of GHRH (10<sup>-7</sup> M) to induce GH secretion. A minimum of 200 pituitary cells were counted in each of three replicate chambers for each treatment to determine the percentage of GH plaque-forming cells. These results are the means and SEM from four independent experiments. Values denoted by different letters are significantly different ( $P < 0.05$ ).

(10<sup>-8</sup> M T<sub>3</sub> for 3 d, 10<sup>-9</sup> and 10<sup>-8</sup> M T<sub>3</sub> for 6 d). Thus, T<sub>3</sub> regulates the induction of somatotroph differentiation by corticosterone in a biphasic, dose-dependent manner.

#### Effect of corticosterone and T<sub>4</sub> alone or in combination on somatotroph abundance

It is well established that circulating T<sub>3</sub> in birds and mammals is derived almost exclusively from the peripheral conversion of T<sub>4</sub> secreted by the thyroid gland. The next experiment was designed to examine the effect of T<sub>4</sub> on somatotroph abundance *in vitro*. E11 pituitary cells were treated with T<sub>3</sub> or T<sub>4</sub> (10<sup>-10</sup> M, 10<sup>-8</sup> M) with and without

corticosterone (10<sup>-9</sup> M) for 3 d (Fig. 2). Similar to results with the RHPA above, corticosterone alone increased proportions of GH-containing cells to 8.9 ± 1.0% *vs.* basal levels of 0.5 ± 1.0%. T<sub>3</sub> at 10<sup>-10</sup> M significantly increased the number of GH-containing cells induced by corticosterone after 3 d of culture (to 12.9 ± 1.1%), but T<sub>3</sub> at 10<sup>-8</sup> M suppressed the effect of corticosterone (to 5.3 ± 1.1%). T<sub>4</sub> alone slightly increased the percentage of GH-containing cells. In contrast to T<sub>3</sub>, T<sub>4</sub> significantly enhanced the effect of corticosterone on somatotroph abundance at both doses. T<sub>4</sub> at 10<sup>-9</sup> M was chosen for the remaining experiments.

The next experiment was designed to determine whether the synergistic interactions with T<sub>4</sub> occurred across a wide concentration range for corticosterone. E11 pituitary cells were treated with corticosterone (10<sup>-11</sup> to 10<sup>-8</sup> M) alone or in combination with T<sub>4</sub> (10<sup>-9</sup> M) for 3 d. Treatment with 10<sup>-10</sup> to 10<sup>-8</sup> M corticosterone significantly increased GH cell abundance in e11 cells, relative to basal (Fig. 3). T<sub>4</sub> augmented the somatotroph abundance induced by all doses of corticosterone tested, indicating that the synergistic effect was not restricted to a limited range of corticosterone levels. However, corticosterone at 10<sup>-10</sup> M in combination with T<sub>4</sub> appeared most enhanced (7.7 ± 1.3%;  $P < 0.01$ ), relative to corticosterone alone (5.5 ± 1.3%). For this reason, this concentration was selected for the remaining trials.

#### Conversion of T<sub>4</sub> to T<sub>3</sub> is necessary for T<sub>4</sub> modulation of somatotroph abundance

The different dose-dependent effects of T<sub>4</sub> and T<sub>3</sub> likely reflect a difference in the mechanism of action of these thyroid hormones. The affinity for its receptor and the biological




FIG. 2. The effect of corticosterone and T<sub>3</sub> or T<sub>4</sub> (TH) on somatotroph abundance determined by ICC. E11 pituitary cells were cultured for 3 d with corticosterone (10<sup>-9</sup> M), T<sub>3</sub> (10<sup>-10</sup> M, 10<sup>-8</sup> M), or T<sub>4</sub> (10<sup>-10</sup> M, 10<sup>-8</sup> M) alone or in combination. Cells were harvested, and the percentage of pituitary cells that contained GH was determined using ICC. A minimum of 200 pituitary cells were counted in each of four replicate wells for each treatment. Data were analyzed by ANOVA, followed by Tukey's Studentized range test. Results shown are the means ± SEM of four independent experiments. Values denoted by different letters are significantly different ( $P < 0.05$ ).

potency of  $T_3$  is 10 times higher than that of  $T_4$  in rat tissues and cultured cell lines derived from the rat and other species (25). It is well known that the type II 5'-deiodinase is responsible for conversion of  $T_4$  to  $T_3$  in the anterior pituitary of adult animals. Iopanoic acid (IOP) is both an inhibitor of all three types of deiodinases and a substrate for the pituitary type II 5'-deiodinase (26, 27). Furthermore, IOP has been shown to block  $T_4$  to  $T_3$  conversion in chick embryos from d

19 of incubation (28). So IOP was used to explore the basis for dose-dependent differences between  $T_4$  and  $T_3$ . Based on preliminary experiments, we determined that  $10^{-6}$  M IOP was not cytotoxic and did not affect corticosterone induction of somatotrophs. We next examined whether the conversion of  $T_4$  to  $T_3$  was necessary for the  $T_4$  effect on somatotroph abundance. Cells from pituitaries of e11 chick embryos were cultured for 3 d in the presence of IOP ( $10^{-6}$  M),  $T_4$  ( $10^{-9}$  M),  $T_3$  ( $10^{-10}$  M), and corticosterone ( $10^{-10}$  M) alone and in combination. Figure 4 presents the results from these trials. Corticosterone increased the somatotroph abundance to  $5.6 \pm 0.9\%$  vs. basal levels of  $2.1 \pm 0.9\%$ .  $T_4$  significantly enhanced the effect of corticosterone on somatotroph abundance (to  $7.5 \pm 0.8\%$ ). No significant effects were detected for IOP and  $T_3$  alone, relative to basal. IOP did not affect the stimulatory effect of corticosterone, as we expected. In contrast, IOP blocked the slight increase in GH cell abundance induced by  $T_4$  and inhibited the added effect of  $T_4$  on corticosterone induction of GH cells (to  $5.5 \pm 0.9\%$ ). However, IOP did not suppress the combined effect of corticosterone and  $T_3$ . Interestingly,  $T_3$  treatment overcame the IOP-induced reduction in the corticosterone/ $T_4$  effect. Thus, conversion of  $T_4$  to  $T_3$  was necessary for  $T_4$  modulation of somatotroph abundance.

## Discussion

It has been demonstrated that somatotroph differentiation does not occur *in vitro* without an extrapituitary signal in rats and chickens. Previously we reported that e16 chicken serum could induce premature somatotroph differentiation, and adrenal corticosterone is the bioactive component of e16 serum (5, 6). We have found that corticosterone is capable of inducing GH cell differentiation *in vitro* and *in vivo* (6, 7). We





FIG. 3. Determination of the range of corticosterone doses for observing synergistic interactions with  $T_4$ . E11 pituitary cells were treated with corticosterone ( $10^{-11}$  to  $10^{-8}$  M) alone or in combination with  $T_4$  ( $10^{-9}$  M) for 3 d. The percentage of GH cells was determined using ICC. Results shown are the means  $\pm$  SEM of four independent experiments. Data were analyzed as in Fig. 2, with values denoted by an asterisk (\*) being significantly different from each other ( $P < 0.05$ ).

FIG. 4. Effect of monodeiodinase inhibition on thyroid hormone effects on somatotroph abundance. E11 pituitary cells were treated with corticosterone ( $10^{-10}$  M),  $T_4$  ( $10^{-9}$  M),  $T_3$  ( $10^{-10}$  M), or IOP ( $10^{-6}$  M) alone or in combination for 3 d. Cells were harvested, and the percentage of GH cells was determined using ICC. Results shown are the means  $\pm$  SEM of four independent experiments. Data were analyzed as in Fig. 2, with values denoted by an asterisk (\*) being significantly different from each other ( $P < 0.05$ ).



have also found that serum concentrations of glucocorticoids increase by e14 (Liu, L., and T. E. Porter, unpublished data). In agreement with our previous work, we found in the present study that short- and long-term exposures to corticosterone increased somatotroph percentages among pituitary cells from e11, an age in which somatotrophs are normally rare.

Thyrotrophs have been identified early in embryonic development, on e6.5 in chickens (29), and functionality of the thyroid axis occurs between e10.5 and 12.5 of development (30). Serum  $T_4$  and  $T_3$  have been detected in chick embryos as early as e9.5 (31, 32). We measured plasma total  $T_4$  and total  $T_3$  from e9 through e14 (Liu, L., and T. E. Porter, unpublished data). Levels of both hormones increased between e11 and e13, with  $T_4$  being twice as abundant as  $T_3$  during this period. Levels of both total  $T_4$  and total  $T_3$  were at levels that were stimulatory to glucocorticoid effects in the present study ( $\sim 10^{-10}$  to  $10^{-9}$  M). Thus, thyroid hormones are present in the systemic circulation from the time of somatotroph differentiation. Thyroid hormones regulate GH gene expression through thyroid hormone receptors. Thyroid hormone receptors are the product of two genes,  $TR\alpha$  and  $TR\beta$ . The chicken  $TR\alpha$  gene produces a single  $TR\alpha$  mRNA, which is translated into two proteins by the use of two alternative start codons, and the  $TR\beta$  gene gives rise to two variant mRNAs,  $TR\beta 0$  and  $TR\beta 2$  (33, 34). The rat GH gene is thought to be a potential target for  $TR\beta 2$  because this receptor has been reported to be highly expressed in rat somatotrophs (35), and it is the predominant TR in  $GH_3$  cell nuclear extract that binds to the GH promoter response element *in vitro* (36). Analysis of expression of  $TR\alpha$  and  $TR\beta$  mRNA during chicken development demonstrated that  $TR\alpha$  mRNA was ubiquitously present from early embryonic stages, whereas  $TR\beta$  mRNA displayed restricted tissue specificity, occurring notably in brain, eye, lung, yolk sac, and kidney, and was subject to striking developmental control (37). However, no report has specifically examined expression in the pituitary.

Multiple studies suggest that thyroid hormones can activate rat GH synthesis and synergistically stimulate GH expression with glucocorticoids in rat pituitary cell lines (16–19). However, thyroid hormones appear to require glucocorticoids to regulate GH gene expression and somatotroph differentiation in fetal animals. Nogami *et al.* (8) found that  $T_3$  increased GH gene expression in cultured fetal rat pituitary glands only in the presence of glucocorticoids. Similarly, although having no effect when given alone,  $T_4$  injections into pregnant rats increased GH and GH mRNA accumulation in the corresponding fetal pituitary glands on d 17 or 18 of gestation, which were induced by dexamethasone in the drinking water (20). Hemming *et al.* (23) demonstrated that GH cells could be induced in cultured fetal rat pituitary primordia with cortisol.  $T_3$ , which was individually ineffective, acted synergistically with cortisol to increase GH cell density in the sectioned tissue. These reports indicate that glucocorticoids and thyroid hormones can act synergistically to induce GH production in the fetal rat pituitary gland. Our present results using dissociated chicken embryonic pituitary cells in culture indicate that the synergistic actions of

glucocorticoids and thyroid hormones occur directly on the somatotroph precursor population to induce an increase in the absolute abundance of GH-producing cells.

To date there have been no reported studies on effects of thyroid hormones on somatotroph abundance during chicken embryonic development. In the present study, we demonstrated that  $T_3$  or  $T_4$  alone had minimal effects on the percentage of somatotrophs in e11 pituitary cell cultures using RHPA and ICC. Thus, thyroid hormones alone could induce GH cell abundance to a small extent. However, induction of somatotrophs by thyroid hormones alone was limited relative to the effects of combined corticosterone and thyroid hormone treatments. It was interesting to find that the corticosterone and  $T_3$  interaction on somatotroph abundance was dose dependent. Similar to the results of Hemming *et al.* using fetal rat pituitary primordia, our data showed that low doses of  $T_3$  in combination with corticosterone could further increase somatotroph proportions above the effect of corticosterone alone. On the other hand, we found that high doses of  $T_3$  inhibited the stimulatory effect of corticosterone on somatotroph differentiation. No similar results were reported in other models.

Compared with  $T_3$ , our data demonstrate that all doses of  $T_4$  can act synergistically with corticosterone to augment corticosterone-induced GH cell differentiation *in vitro*. The primary hormone secreted by the thyroid gland is  $T_4$ . Most circulating  $T_3$  is produced by monodeiodination of  $T_4$  in liver, kidney, brain, pituitary, and other peripheral tissues. Type I or II deiodinase activity has been demonstrated in several tissues of fetal rats (38, 39) and embryonic chickens (40), including the pituitary. In cultured  $GH_3$  cells, addition of ipodate, a compound closely related to IOP, caused inhibition of  $T_4$  5'-deiodination and prevented stimulation of GH synthesis by  $T_4$  (41). Our data showed that IOP did not affect somatotroph differentiation induced by corticosterone alone or by the corticosterone/ $T_3$  combination. In contrast, IOP blocked the slightly stimulatory effect of  $T_4$  and the combined effect of corticosterone/ $T_4$ . Addition of  $T_3$  reversed the inhibition effect of IOP. Thus, conversion of  $T_4$  to  $T_3$  was necessary for the effect of  $T_4$  on corticosterone induction of somatotrophs, suggesting that  $T_3$  plays the primary role in thyroid hormone regulation of somatotroph differentiation during development. The results of our IOP experiments may also explain the different dose-dependent effects between  $T_4$  and  $T_3$ . Our data indicate that low levels of  $T_3$  acted synergistically with corticosterone to augment corticosterone-induced GH cell abundance. In contrast, the high dose of  $T_3$  ( $10^{-8}$  M), which is higher than its physiological level at that age, inhibited the stimulatory effect of corticosterone on somatotroph differentiation. These findings suggest that the capacity of the anterior pituitary to convert  $T_4$  to  $T_3$  is likely limiting at this age of chicken embryonic development, thereby restricting the effective local concentration of  $T_3$  to low levels, even in the presence of high  $T_4$ . If pituitary  $T_4$  monodeiodinase activity is limiting, this would allow for stimulatory effects of thyroid hormones on somatotrophs and limit potentially inhibitory effects of higher circulating  $T_4$  concentrations.

## Acknowledgments

Received February 3, 2003. Accepted May 19, 2003.

Address all correspondence and requests for reprints to: Dr. Tom E. Porter, Department of Animal and Avian Sciences, University of Maryland, College Park, Maryland 20742. E-mail: tp44@umail.umd.edu.

This work was supported by competitive Grant 00-35206-9463 (to T.E.P.) from the United States Department of Agriculture National Research Initiative program.

## References

- Treier M, Gleiberman AS, O'Connell SM, Szeto DP, McMahon JA, McMahon AP, Rosenfeld MG 1998 Multistep signaling requirements for pituitary organogenesis *in vivo*. *Genes Dev* 12:1691–1704
- Treier M, Rosenfeld MG 1996 The hypothalamic-pituitary axis: co-development of two organs. *Curr Opin Cell Biol* 8:833–843
- Li S, Crenshaw III EB, Rawson EJ, Simmons DM, Swanson LW, Rosenfeld MG 1990 Dwarf locus mutants, which lack three pituitary cell types, result from mutations in the POU domain gene, Pit-1. *Nature* 347:528–533
- Porter TE, Couger GS, Dean CE, Hargis BM 1995 Ontogeny of growth hormone (GH)-secreting cells during chicken embryonic development: initial somatotrophs are responsive to GH-releasing hormone. *Endocrinology* 136:1850–1856
- Porter TE, Couger GS, Morpurgo B 1995 Evidence that somatotroph differentiation during chicken embryonic development is stimulated by a blood borne signal. *Endocrinology* 136:3721–3728
- Morpurgo B, Dean CE, Porter TE 1997 Identification of the blood-borne somatotroph-differentiating factor during chicken embryonic development. *Endocrinology* 138:4530–4535
- Dean CE, Morpurgo B, Porter TE 1999 Induction of somatotroph differentiation *in vivo* by corticosterone administration during chick embryonic development. *Endocrine* 11:151–156
- Nogami H, Inoue K, Kawamura K 1997 Involvement of glucocorticoid-induced factor(s) in the stimulation of growth hormone expression in the fetal rat pituitary gland *in vitro*. *Endocrinology* 138:1810–1815
- Nogami H, Tachibana T 1993 Dexamethasone induces advanced growth hormone expression in the fetal rat pituitary gland *in vivo*. *Endocrinology* 132:517–523
- Paek I, Axel R 1987 Glucocorticoids enhance stability of human growth hormone mRNA. *Mol Cell Biol* 7:1496–1507
- Porter TE, Dean CE, Piper MM, Medvedev KL, Ghavam S, Sandor J 2001 Somatotroph recruitment by glucocorticoids involves induction of growth hormone gene expression and secretagogue responsiveness. *J Endocrinol* 169:499–509
- Scanlon MF, Issa BG, Dieguez C 1996 Regulation of growth hormone secretion. *Horm Res* 46:149–154
- Muller EE, Locatelli V, Cocchi D 1999 Neuroendocrine control of growth hormone secretion. *Physiol Rev* 79:511–607
- Chomczynski P, Soszynski PA, Frohman LA 1993 Stimulatory effect of thyroid hormone on growth hormone gene expression in a human pituitary cell line. *J Clin Endocrinol Metab* 77:281–285
- Hervas F, Morreale de Escobar G, Escobar Del Rey F 1975 Rapid effects of single small doses of L-thyroxine and triiodo-L-thyronine on growth hormone, as studied in the rat by radioimmunoassay. *Endocrinology* 97:91–101
- Shapiro LE, Samuels HH, Yaffe BM 1978 Thyroid and glucocorticoid hormones synergistically control growth hormone mRNA in cultured GH1 cells. *Proc Natl Acad Sci USA* 75:45–49
- Dobner PR, Kawasaki ES, Yu LY, Bancroft FC 1981 Thyroid or glucocorticoid hormone induces pre-growth-hormone mRNA and its probable nuclear precursor in rat pituitary cells. *Proc Natl Acad Sci USA* 78:2230–2234
- Spindler SR, Mellon SH, Baxter JD 1982 Growth hormone gene transcription is regulated by thyroid and glucocorticoid hormones in cultured rat pituitary tumor cells. *J Biol Chem* 257:11627–11632
- Yaffe BM, Samuels HH 1984 Hormonal regulation of the growth hormone gene. Relationship of the rate of transcription to the level of nuclear thyroid hormone-receptor complexes. *J Biol Chem* 259:6284–6291
- Nogami H, Yokose T, Tachibana T 1995 Regulation of growth hormone expression in fetal rat pituitary gland by thyroid or glucocorticoid hormone. *Am J Physiol* 268:E262–E267
- Scanes CG, Duyka DR, Lauterio TJ, Bowen SJ, Huybrechts LM, Bacon WL, King DB 1986 Effect of chicken growth hormone, triiodothyronine and hypophysectomy in growing domestic fowl. *Growth* 50:12–31
- Denver RJ, Harvey S 1991 Thyroidal inhibition of chicken pituitary growth hormone: alterations in secretion and accumulation of newly synthesized hormone. *J Endocrinol* 131:39–48
- Hemming FJ, Aubert ML, Dubois PM 1988 Differentiation of fetal rat somatotropes *in vitro*: Effects of cortisol, 3, 5, 3'-triiodothyronine, and glucagons, a light microscopic and radioimmunological study. *Endocrinology* 123:1230–1236
- Dean CE, Porter TE 1999 Regulation of somatotroph differentiation and growth hormone secretion by corticosterone and GH-releasing hormone during embryonic development. *Endocrinology* 140:1104–1110
- Samuels HH, Forman BM, Horowitz ZD, Ye ZS 1989 Regulation of gene expression by thyroid hormone. *Annu Rev Physiol* 51:623–639
- Lam SK, Harvey S 1985 *In vitro* conversion of thyroxine to triiodothyronine by chicken hepatic 5'-deiodinase: kinetic studies. *J Endocr* 110:441–446
- Kaplan MM 1984 The role of thyroid hormone deiodination in the regulation of hypothalamo-pituitary function. *Neuroendocrinology* 38:254–260
- Decuypere E, Kühn ER, Clijmans B, Nouwen EJ, Michels H 1982 Effect of blocking T<sub>4</sub>-monodeiodination on hatching in chickens. *Poultry Sci* 61:1194–1201
- Thommes RC, Fitzsimons EJ, Davis M, Woods JE 1992 Immunocytochemical demonstration of T<sub>4</sub> content and TSH-binding by cells of the thyroid of the developing chick embryo. *Gen Comp Endocrinol* 85:79–85
- Thommes RC 1987 Ontogenesis of thyroid function and regulation in the developing chick embryo. *J Exp Zool Suppl* 1:273–279
- Thommes RC, Vieth RL, Lévassieur S 1977 The effects of hypophysectomy by means of surgical decapitation on thyroid function in the developing embryo. *Gen Comp Endocrinol* 31:29–36
- Thommes RC, Hylka VW 1977 Plasma iodothyronine in the embryonic and immediate post-hatch chick. *Gen Comp Endocrinol* 32:417–422
- Forrest D, Hallböök F, Persson H, Vennström B 1991 Distinct function for thyroid hormone receptors  $\alpha$  and  $\beta$  in brain development indicated by differential expression of receptor genes. *EMBO J* 10:269–275
- Sjöberg M, Vennström B, Forrest D 1992 Thyroid hormone receptors in chick retinal development: differential expression of mRNAs for  $\alpha$  and N-terminal variant  $\beta$  receptors. *Development* 114:39–47
- Childs, GV, Taub K, Jones KE, Chin WW 1991 Triiodothyronine receptor  $\beta$ -2 messenger ribonucleic acid expression by somatotropes and thyrotropes: effect of propylthiouracil-induced hypothyroidism in rats. *Endocrinology* 129:2767–2773
- Suen C-S, Chin WW 1993 *In vitro* transcriptional studies of the roles of thyroid hormone (T<sub>3</sub>) response elements and minimal promoters in T<sub>3</sub> stimulated gene transcription. *J Biol Chem* 269:1314–1322
- Forrest D, Sjöberg M, Vennström B 1990 Contrasting developmental and tissue-specific expression of  $\alpha$  and  $\beta$  thyroid hormone receptor genes. *EMBO J* 9:1519–1528
- Bates JM, Germain DL, Galton VA 1999 Expression profiles of the three iodothyronine deiodinases, D1, D2, and D3, in the developing rat. *Endocrinology* 140:844–851
- Rodriguez-Garcia M, Jolin T, Santos A, Perez-Castillo A 1995 Effect of perinatal hypothyroidism on the developmental regulation of rat pituitary growth hormone and thyrotropin genes. *Endocrinology* 136:4339–4359
- Van der Geyten S, Van den Eynde I, Segers IB, Kuhn ER, Darras VM 2002 Differential expression of iodothyronine deiodinases in chicken tissues during the last week of embryonic development. *Gen Comp Endocrinol* 128:65–73
- Melmed S, Nelson M, Kaplowitz N, Yamada T, Hershman JM 1981 Glutathione-dependent thyroxine 5'-monodeiodination modulates growth hormone production by cultured nonthyrotropic rat pituitary cells. *Endocrinology* 108:970–976